



CLINICAL REVIEW

Attention deficit hyperactivity disorder and sleep disordered breathing in pediatric populations: A meta-analysis



Karim Sedky^{a,*}, David S. Bennett^b, Karen S. Carvalho^b

^a Cooper Hospital Rowan University, Camden, NJ, USA

^b Drexel University College of Medicine, Philadelphia, PA, USA

ARTICLE INFO

Article history:

Received 13 July 2013

Received in revised form

25 November 2013

Accepted 15 December 2013

Available online 24 December 2013

Keywords:

Sleep disordered breathing

Obstructive sleep apnea

Attention deficit hyperactivity disorder

Meta-analysis

SUMMARY

A relationship between attention deficit hyperactivity disorder (ADHD) and sleep disordered breathing (SDB) in children and adolescents has been suggested by some authors. Yet, this topic remains highly controversial in the literature.

A meta-analysis was conducted in order to examine the extent of relationship between SDB and ADHD symptoms in pediatric populations and whether there are differences in ADHD symptoms pre- versus post-adenotonsillectomy in pediatric populations.

PubMed/Medline, PsychInfo and Cochrane databases were searched using the key words “attention deficit hyperactivity disorder” or “ADHD” and “obstructive sleep apnea” or “OSA” or “sleep disordered breathing” (SDB) or “SDB”. English language publications through September 2012 were surveyed. Meta-analysis was conducted to assess the relationship between SDB and ADHD symptoms in the first part of the study, and the extent of change in ADHD symptoms before and after adenotonsillectomy in the second part.

Eighteen studies satisfied the inclusion criteria for the first part of the study. This represented 1113 children in the clinical group (874 diagnosed with SDB who were examined for ADHD symptoms; 239 diagnosed with ADHD who were examined for SDB) and 1405 in the control-group. Findings indicate that there is a medium relationship between ADHD symptoms and SDB (Hedges' $g = 0.57$, 95% confidence interval: 0.36–0.78; $p = 0.000001$). A high apnea hypopnea index (AHI) cutoff was associated with lower effect sizes, while child age, gender and body mass index did not moderate the relationship between SDB and ADHD. Study quality was associated with larger effect sizes. In the second part of the study, twelve studies were identified assessing pre- versus post-surgery ADHD symptoms. Hedges' g was 0.43 (95% confidence interval = 0.30–0.55; $p < 0.001$; $N = 529$) suggesting a medium effect, as adenotonsillectomy was associated with decreased ADHD symptoms at 2–13 months post-surgery.

The findings of this meta-analysis suggest that ADHD symptoms are related to SDB and improve after adenotonsillectomy. Therefore, patients with ADHD symptomatology should receive SDB screening. Treatment of comorbid SDB should be considered before medicating the ADHD symptoms if present.

© 2013 Published by Elsevier Ltd.

Introduction

Attention deficit hyperactivity disorder (ADHD) is a neuropsychiatric syndrome affecting 3–16% of school children with world prevalence of 5.29% [1]. It is characterized by inattention, hyperactivity, and/or impulsivity, occurring in at least two different

settings, and often leads to academic, social, and occupational dysfunction [2]. Although the diagnosis of ADHD is best made through a clinical interview, rating scales completed by teachers and parents have been utilized and shown to be valid [3]. The variation of diagnostic criteria for ADHD over the years (i.e., using different versions of Diagnostic and Statistical Manual for Mental Disorders or International Classification of Diseases criteria), the source of informant (e.g., parent, teacher, or clinical), and the reliability of rating scales to aid in diagnosis render this diagnosis difficult in some instances [4]. Other psychiatric disorders can also mimic ADHD symptomatology (e.g., anxiety), as may sleep disorders, thus complicating the diagnosis of ADHD.

* Corresponding author. Cooper Hospital – Rowan University, Department of Psychiatry, 3 Cooper Plaza, Camden, NJ 08103, USA. Tel.: +1 (856) 757 7799; fax: +1 (856) 757 9651.

E-mail address: sedky66@hotmail.com (K. Sedky).

Sleep disordered breathing (SDB) in children is a disorder varying from primary snoring (PS) at the least severe end of the spectrum to upper airway resistance syndrome (UARS), obstructive hypoventilation, and obstructive sleep apnea (OSA) in more severe cases. UARS is characterized by partial upper airway obstruction without oxygen desaturation, and is also associated with respiratory event related arousals and more negative esophageal pressure. OSA diagnosis depends on the apnea/hypopnea index (AHI), which is the average number of respiratory apneas and hypopneas per hour of sleep. Grading of the sleep apnea syndrome in children differs from adults; although recommended cutoffs have varied over time, it is generally agreed that an AHI ≥ 1 /h is abnormal [5]. Mild cases of OSA range from 1 to 5 per hour, with moderate cases ranging from 5 to 10 per hour, and severe cases above 10 per hour. Although the recommendations set by the American Academy of Sleep Medicine (AASM) in 2007 specified the criteria of using two missed breaths equivalent duration to diagnose respiratory events, the AHI cutoff limit to diagnosing OSA remains controversial [6].

Several studies suggest an elevated incidence of SDB, ranging from 25 to 57%, among children and adolescents diagnosed with ADHD symptoms [7–13]. Other studies, however, question this relationship as they find no such elevated incidence of ADHD symptoms associated with SDB [14–17]. Small sample sizes, lack of polysomnographic (PSG) diagnosis of SDB, variability between studies on AHI cutoffs used to diagnose SDB (e.g., some used a cutoff of > 5 per hour, while others used > 1 per hour), and variability in whether a clinical diagnosis of ADHD or an ADHD rating scale was used to assess ADHD symptoms might explain some of the inconsistency in outcomes.

SDB during sleep is usually associated with lower oxygen saturations. This intermittent hypoxia/hypercapnia, with consequent sleep interruption might have an impact on brain development and affect cognitive function [18], leading to inattention [19]. Children diagnosed with SDB have been found to have elevated levels of inflammatory cytokines (C-reactive protein and interleukin-6), which also might contribute to poor cognitive function, including poor attention span [20,21].

Due to the inconsistent relationship between SDB and ADHD symptoms, further research is needed to address the extent to which they are related. An improved understanding of this relationship is important as it may have clinical implications. For example, children suffering from mild OSA and ADHD, treated with adenotonsillectomy (AT; the main treatment for SDB in children) had comparable improvements in ADHD symptoms to those treated with methylphenidate in one study [22]. This finding suggests that at least for some children with comorbid SDB, ADHD symptoms can improve with AT surgery without the need for long-term drug treatment. Although possible, it remains unclear whether AT surgery would lower the medication dosage needed for those with residual ADHD symptoms. Early identification and treatment for SDB might prevent permanent life-long disability and financial loss. Using meta-analysis, the current review examines the relation between ADHD and SDB among children and adolescents.

Methods

Study selection

A PubMed/Medline, PsychInfo and Cochrane library search was performed using the terms “attention deficit hyperactivity disorder” or “ADHD”, “attention problems” and “obstructive sleep apnea” or “OSA” or “sleep disordered breathing” or “SDB” crossed by “child”, “children” and “adolescent”. Google Scholar was also used to facilitate the search results. References cited in these articles

were also examined and included if they met study criteria. English language studies through September 2012 were examined.

Inclusion and exclusion criteria

Since this study focuses on the pediatric population, only patients aged from 0 to 18 years old were included. No study was included if the oldest individual was older than 18 years of age.

Although there are several proposed methods to diagnose SDB including the pediatric sleep questionnaire (PSQ), the gold standard diagnostic test remains polysomnography (PSG) [23]. Thus, studies having PSG performed in both clinical and control groups were required for inclusion in the first part of the study. Parent and/or teacher ADHD symptom rating scales or clinician-based ADHD diagnosis were used to assess ADHD symptoms in these subjects. Studies were only included if there was a control group. Authors of studies with missing information were contacted; these studies were only included if sufficient information to compute an effect size (ES) was provided by the researcher. In the second part of the meta-analysis, PSG was required to be done both before and 1–13 months after adenotonsillectomy. ADHD symptomatology was also assessed during the same time frame at both baseline and post-surgery.

Identification of appropriate studies was conducted by the first author (K.S.) and was reviewed by the second author (D.B.). For the first part of the meta-analysis, 48 published studies were initially identified, with ten excluded due to lack of the use of PSG to detect SDB at least in one group [8,9,24–31]. Five others lacked a control group and were excluded [14,32–35]. Another five were excluded for the following reasons: one study selected only for children who had both OSA and ADHD in the clinical group [22], one study did not specify the relationship between AHI and ADHD symptoms [36], one had PSG done but did not present AHI results [10], and two used excessive daytime sleepiness rather than SDB; this latter symptom can originate from a variety of causes other than SDB and was thus excluded [37,38]. In yet another study, the authors used an unvalidated, two-item ADHD scale to compare the two groups; thus, this study was excluded [39]. In addition, data from one study was reported in four different publications; thus, data were used from these four studies [11,40–42] but reported as one study [11]. An additional four studies also appeared to have patient overlap [13,43–45]. After contacting two authors, the extent of patient overlap could not be resolved as one author suggested common patients among two studies [43,44], while another suggested common patients among each of the four. Thus, two different analyses were conducted: one including three out of the four studies and another only including the study with the greatest number of patients in both the OSA and control groups. To minimize positive selection bias of publication, we searched for unpublished studies and identified two dissertations [46,47], both of which had a related publication already identified, one of which was included in the meta-analysis [48]. The other was excluded due to the lack of PSG. In the second part of the study, 12 studies satisfied the selection criteria and were included for meta-analysis [11,13,22,30,33,41,44,45,49–55].

Data extraction

Data from the remaining 20 studies were coded by one author (K.S.) and reviewed by the second (D.B.) for accuracy in the initial part of the study [11–13,15–17,40,41,44,45,48,49,56–65]. Coded information included: sample size, type of clinical group, age, gender, body mass index (BMI), AHI cutoff, and study quality for both clinical and control groups. In the second part, pre- and post-AT ADHD symptoms levels were recorded and differentiated.

Table 1

Studies examining the relation between ADHD symptoms and SDB.

Study Name, year	Variable in clinical group	ADHD and/or OSA depended on	# Clinical subjects	# Control subjects	Mean/SD (age clinical group, y)	Mean/SD (age control group, y)	BMI clinical (kg/m ²)	BMI control (kg/m ²)	Male% clinical group	Male% control group
Barnes et al., 2012 [64]	OSA	CBCL	14	14	6.2 ± 1.81 (4–8)	6.2 ± 1.3 (4–8)	19.4 ± 4.79	17.1 ± 2.5		
Beebe et al., 2010 [65]	SDB	BASC	126	37	13.7 ± 1.9 (10–16.9)	13 ± 2 (10–16.9)	39 ± 7.5	35.9 ± 6.5		
Blunden et al., 2000 [56]	PS + one case of OSA	CBCL	16	16	7.2 ± 1.6 (5–11)	7.7 ± 1.6 (5–11)			43.8	43.8
Bourke et al., 2011 [57]	PS/OSA	CBCL/BRIEF	99	35	9.6 ± 1.5 (7–12)	9.5 ± 1.8			59.3	48.6
Chervin et al., 2001 [15] ^a	SDB	CPRS (hyperactivity index t-score); Clinician Dx	59	54	9.6 ± 3.7 (2–18)	10.1 ± 4.4			66	63
Chervin et al., 2006 [40]	AT/OSA	CPRS, CSI, CPT	40	27	7.8 ± 1.8	9.3 ± 2		18.9 ± 2.9	52.5	70.4
Cooper et al., 2004 [17]	ADHD	CBCL, CPRS, CPT, Clinician Dx	18	20	10.5 ± 3 (6–16)	10 ± 3.9 (4–16)			83.3	55
Galland et al., 2011 [16]	ADHD	Clinician Dx	28	28	10.1 (6–12)	10.2 (6–12)	18.7 ± 3.9	18.7 ± 3.5	78.6	78.6
Golan et al., 2004 [59]	ADHD/OSA	Clinician Dx, OSA Dx according to AHI level with cutoff AHI ≥ 1/h	34	32	12.4 ± 4.6	12 ± 3.6	20.4 ± 3.9	18.6 ± 2.2	76.5	65.6
Gottlieb et al., 2004 [58]	OSA	CPRS	61	144	5	5			51.7	
Huang et al., 2004 [12]	ADHD	CBCL, Clinician Dx	88	27	8.5 ± 1.9 (6–12)	9 ± 2 (6–12)	18.3 ± 3.7	20.0 ± 3.7	87.5	77.8
Kaemingk et al., 2003 [48] ^a	OSA	CPRS-R, Home sleep study	77	69	8.3 ± 1.8 (6–12)	8.4 ± 0.6			54.4	
Landau et al., 2012 [60]	OSA	CBCL, C-TRF	45	13	3.8 ± 9.2 (2.5–5)	4.1 ± 0.7	15.7 ± 2	16.4 ± 2	73.3	46.2
Lewin et al., 2002 [49]	Mild/Sev OSA	CBCL	21	10	7.4 ± 2.7 (4–12)	6.9 ± 3.6 (4–9)	18.4 ± 4.9	16.3 ± 3.6	50	80
Miano et al., 2011 [61]	PS/OSA	ADHD rating scale	44	60	8.8 ± 1.9	9.8 ± 2.4	18.6 ± 3.6	18.3 ± 1.8	60.5	36.7
O'Brien et al., 2003 [13] ^a	Mild ADHD	CBCL, CPRS	71	39	6.5 (5–7)	6.5 (5–7)				
O'Brien et al., 2003 [44]	ADHD	Clinician Dx	100	49	7.2 ± 0.9	6.7 ± 0.4			66	44
O'Brien et al., 2004 [45]	SDB	CPRS, CBCL	35	35	6.7 ± 0.6 (5–7)	6.7 ± 0.5 (5–7)	19.8 ± 4.3	17.7 ± 3.5	49	49
Rosen et al., 2004 [62]	OSA	CBCL, CPRS	40	667	9.3 ± 0.9 (8–11)	9.5 ± 0.8	19 ± 4.7	17.9 ± 3.5	42.5	50.2
Ting et al., 2011 [63]	OSA	CBCL, TRF	128	10	10.2 ± 1.2	10.1 ± 0.9	18.4 ± 3.2	17.1 ± 0.8	63.3	40

ADHD: Attention Deficit Hyperactivity Disorder; BRIEF: Behavior Rating Inventory of Executive Function; CBCL: Child Behavior Check List; CPRS: Conner's parent rating scale; CPT: Auditory continuous performance test; CRPS-R: Conners' Parent Rating Scales-Revised; CSI: child symptom inventory; CTRS: Conner's teacher rating scale; C-TRF: caregiver teacher report; NEPSY: neuropsychological assessment; OSA: obstructive sleep apnea; PS: primary snoring; SD: Standard Deviation; SDB: sleep disordered breathing includes PS and OSA.

^a Studies that used AHI cutoff ≥5/h to diagnose OSA.

Data analysis

Data from the studies were analyzed using the Comprehensive Meta-Analysis, version 2.0 (Biostat, Englewood, NJ) software program. Hedges' *g* is an effect size statistic that represents the standardized difference between two groups. It was computed using a random effects model, with 95% confidence intervals (CI) reported. The results were plotted on a Forrest graph to provide an overview of the results. Homogeneity between studies was tested using *Q*, while *I*² index assessed the degree of their heterogeneity, estimating the total variation across studies caused by heterogeneity rather than chance. A value of *I*² around 25% suggests low heterogeneity, 50% as moderate, while 75% suggests high heterogeneity. Publication bias was evaluated by examining the relationship between standard errors of the means and the effect sizes and was estimated by the Kendall's tau value. To assess the potential moderator effects of age, gender, BMI, study quality, and AHI cutoff, the mean for each variable was computed for each study and weighted by sample size, and then analyzed in Statistical Product and Service Solutions (SPSS)[®] using linear regression analyses using inverse variances as weights. Study quality was assessed using the Newcastle–Ottawa Scale for case–control studies [66]. This scale has eight items divided into three different categories: selection, comparability and exposure. Each item is graded with a point, if it satisfies certain level except comparability which can have up to two points. Thus, a total score for study quality can range from zero to a maximum of nine points.

Results

Question 1: Is there a relationship between SDB and ADHD symptomatology?

Eighteen studies were included in the analysis constituting a total of 1113 children in the clinical group (mean number per study is 62) that were compared to 1405 in the control group (with mean of 78 individuals in each study; see Table 1). The mean unweighted age for the clinical group was 8.37 years (SD = 2.32) and 8.54 years (SD = 2.23) for the controls. The clinical groups consisted of two: one in which children with SDB (874 individuals) were evaluated with ADHD rating scales and compared to controls; the second in which children diagnosed with ADHD (239) were administered PSG and compared to controls.

Aggregate effect sizes and homogeneity tests

The overall ES (Hedges' *g*) for the eighteen studies assessing the relation between ADHD symptoms and SDB was 0.57 (95% CI = 0.36–0.78; *p* = 0.00001; see Fig. 1). After the inclusion of the other two studies [44,45], raising the overall number of studies to twenty, the overall ES for the relation between ADHD symptoms and SDB was 0.53 (95% CI = 0.34–0.72; *p* = 0.00001). Thus, it does not appear that the addition of these two studies would have changed the results significantly. Of note, one of the excluded studies [39] used an unvalidated ADHD measure but had an ES of 0.67 (CI = 0.44–0.90; *p* ≤ 0.001) similar to 0.57, suggesting that exclusion of this study also did not greatly affect the overall

reported ES. Testing for heterogeneity, the Q value with 17 d.f. was 71.30, signifying considerable heterogeneity across studies, while I^2 was 76.16, $p < 0.001$, confirming this high degree of heterogeneity. Therefore, we tested several potential moderators as possible sources of this heterogeneity. In addition, Kendall's tau = 0.25; $p = 0.15$; this suggests that the standard errors of the means and the effect sizes were independent, suggesting that there was no significant publication bias.

Moderator analyses

Utilizing a mean weighted ES among the eighteen studies to assess for moderator effects, significant moderation was found for AHI scores such that studies using a higher AHI cutoff had lower ESs (beta = -0.61 , $p = 0.007$). Among the five studies using a high AHI cutoff (i.e., AHI of ≥ 4 –5/h), the ES was 0.26 (95% CI = 0.08–0.44; $p = 0.004$) [13,15,48,58,62]. This was compared to the rest of the studies using a lower AHI cutoff (i.e., AHI ≥ 1), which had a higher ES of 0.72 (95% CI = 0.46–0.99; $p = 0.0000001$). A significant moderation effect also was found for study quality as studies of better quality had larger ESs (beta = 0.48, $p = 0.045$). No significant moderation was found for age (beta = 0.07, $p = 0.80$), gender (beta = -0.33 , $p = 0.25$) or BMI (beta = 0.34, $p = 0.42$). We also examined whether ESs were greater for inattention than for hyperactivity symptoms, using four studies that assessed both sets of symptoms [48,61,62,65]. The mean ES for hyperactivity symptoms with OSA was 0.54 (95% CI = 0.05–1.03; $p = 0.03$), while the mean ES for inattention symptoms was similar at 0.50 (95% CI = 0.03–0.98; $p = 0.038$). These subscales were not significantly different and suggest that the type of ADHD symptoms does not moderate the relationship between ADHD and OSA.

Question 2: Is there any difference between ADHD symptomatology pre- versus post-AT?

In the second part of the study, which involved 529 individuals, ADHD symptoms were compared before surgery versus post-AT among children with SDB (see Table 2). Hedges' g was 0.43 (95%

CI = 0.30–0.55; $p < 0.001$), indicating a medium improvement in ADHD symptoms from pre- to post-surgery (see Fig. 2). Testing for heterogeneity, the Q value with 11 d.f. was 10.03, signifying minimal heterogeneity across studies.

Discussion and conclusion

Pediatric populations suffering from SDB are at increased risk of presenting with symptoms of ADHD, including inattention and hyperactivity. The ES of 0.57 found in the current meta-analysis suggests that a medium relationship exists between SDB and ADHD symptoms. Moreover, a medium improvement was found in ADHD symptoms following adenotonsillectomy (ES = 0.43).

Substantial heterogeneity was observed across studies, including across prior meta-analyses. Of the two prior meta-analyses examining SDB and ADHD symptoms, which were based on much smaller samples of children, one found a medium ES similar to ours meta-analysis [67]. Because the authors used strict criteria in defining group membership (e.g., they excluded studies in which children had comorbid psychiatric diagnoses aside from oppositional behavior, and also excluded those whose ADHD was treated with medication), only three studies that were also included in our meta-analysis were included in their review. Their findings, however, were consistent with those of the current, larger meta-analysis as they reported an overall effect size of 0.52 (CI = 0.23–0.81) across the three studies. In the second meta-analysis differentiating AHI in children with ADHD versus controls, a smaller ES (Cohen's d) of 0.10 was found (95% CI = -0.40 to 0.60), suggesting no relationship between AHI scores in OSA patients and ADHD symptoms [4]. Their meta-analysis, however, was published in 2006 and limited to only four studies totaling 227 children in the ADHD group versus 132 in the control group. Since these two prior meta-analyses were conducted earlier, they were unable to include a number of recent publications addressing the relationship between SDB and ADHD symptoms.

In the large prospective TuCASA study of children 6–12 years of age [68], an overall ES of 0.31 was found between respiratory disturbance index (high vs. low) and parent ratings of children's




Fig. 1. Randomized effect size (Hedges' g) for the relationship between attention deficit hyperactivity disorder (ADHD) symptoms and sleep disordered breathing (SDB).

Table 2

Studies examining ADHD symptoms both before and after adenotonsillectomy.

Study Name, year	ADHD scale used	# Pre-AT	# Post-AT	Mean/SD age (age range)	Duration between assessment (months)	Pre-AT (#/h)	Post-AT AHI (#/h)	Cut-off/h	Male%
Chervin et al., 2012 [55]	CPRS & CSI	81	81	7.8 ± 2.8	7.2 ± 0.8	RDI = 7.5 ± 11.1	2.1 ± 1.8		54.32
Dillon et al., 2007 [11]	DBDRS-combined	40	39		13.0 ± 1.4				
Galland et al., 2006 [30]	BASC-P	61	61	7.0 ± 2.0 (4–11)	3	AHI = 3.0 ± 2.6		AHI ≥ 1.5	57.38
Giordani et al., 2012 [41]	CPRS	40	39	7.8 ± 1.8	12	OAI = 1.5 ± 0.45	0.18 ± 0.45	OAI ≥ 0.5	52.5
Huang et al., 2007 [22]	CBCL-P & ADHD-RS	25	25	8.08 ± 1.28 (6–12)		AHI = 3.3 ± 1.1	0.89 ± 0.63	AHI ≥ 1	92.0
Lewin et al., 2002 [49]	CBCL	7	7	(4–12)	6–18	RDI = 17.4 ± 9.9		RDI ≥ 10	
Li et al., 2006 [33]	CBCL	40	40	8.4 ± 1.6	6	AHI = 10.6 ± 11.1	1.7 ± 2.1	AHI ≥ 1	90.0
Mitchell et al., 2005 [50]	BASC	52	52	7.1 (5–88)	4.4	AHI = 16.2		AHI ≥ 5	56
Mitchell et al., 2006 [51]	BASC	23	23	7.2 (2.5–14.8)	2.4	AHI = 14.1		AHI ≥ 5	65
Mitchell et al., 2007 [52]	BASC	40	40	7.07	6	AHI = 15.9		AHI ≥ 5	55
Mitchell et al., 2009 [53]	BASC	89	89	(3–17.1)	3–6	AHI = 16.3	4.7	AHI ≥ 2	60.67
Tran et al., 2005 [54]	BASC	31	31	5.8 ± 2.5 (2–11.5)	5.4 ± 1.9			RDI ≥ 5 or AHI ≥ 1	60

ADHD: Attention deficit hyperactivity disorder; ADHD-RS: ADHD rating scale; AHI: apnea hypopnea index; AT: adenotonsillectomy; BASC-P: behavioral assessment scales for children (parent); CBCL: child behavior checklist; CPRS: Conner's parent rating scale; CSI: child symptoms inventory; DBDRS: disruptive behavior disorders rating scale; OAI: obstructive apnea index; RDI: respiratory disturbance index.

ADHD symptoms. This study was not included in the current meta-analysis because a somewhat arbitrary cut-off was made between what constituted a high vs. low RDI score (i.e., “high” represented the highest 15%, while “low” represented the lowest 85%), but the study's ES, while somewhat lower than that found in our meta-analysis, was significant and indicated a small to medium relationship between SDB and ADHD symptoms. Guidelines for diagnosing pediatric OSA were published in 2007, after this study was completed [6].

Several potential moderators of the relation between ADHD symptoms and SDB were examined in the current meta-analysis. Study AHI cutoff was found to moderate the ADHD–SDB relationship such that studies using lower AHI cutoffs actually had higher ESs. There are five studies utilizing an AHI cutoff of ≥ 4 –5/h to diagnose OSA; two of these studies did not find a relation between ADHD symptoms and SDB/OSA. This suggests that the milder form of OSA might actually be more closely related to ADHD symptoms, although the reason for this finding is unclear. Study quality also was found to be a significant moderator as higher quality studies had larger ESs, suggesting that the ADHD–SDB relationship is not an artifact of poorer quality studies.

Adenotonsillectomy remains the main treatment for OSA in the pediatric population with enlarged lymphoid tissue. A recent

relatively large, randomized trial of 210 children, age 5–9 y was not included in the current meta-analysis because researchers compared the caregiver and the teacher ratings for children using a global rating scale of psychopathology as opposed to one focusing specifically on ADHD symptoms [69]. Similar to our findings for ADHD symptoms, however, a mild to moderate decrease was observed in the global rating scale as assessed by both groups (caregiver ES = 0.28; teacher's ES = 0.37), raising the possibility that symptoms other than ADHD symptoms might also be improved following adenotonsillectomy. Consistent with this hypothesis, a recent meta-analysis found depressive symptoms to decrease significantly among children and adolescents with SDB who underwent adenotonsillectomy [70].

There are several possible explanations for the relationship between SDB and ADHD symptoms. SDB syndrome is associated with lower oxygen saturations and hypercapnia overnight, oxidative stress, increased free radicals, and/or release of inflammatory cytokines (e.g., proinflammatory cytokines interleukin-6 and tumor necrotizing factor alpha) leading to neurological dysfunction particularly involving certain brain areas (e.g., prefrontal cortex [20,21,71]). Cortical dysfunction is associated with cognitive and behavior dyscontrol and can consequently lead to inattention,




Fig. 2. Pre- versus post-adenotonsillectomy (AT) as compared by the attention deficit hyperactivity disorder (ADHD) symptom level. Group-A signifies pre-AT while group-B is post-AT.

hyperactivity or impulsivity symptoms [72]. SDB also frequently leads to sleep fragmentation, thus, rendering the child exhausted and cognitively impaired during the day [47]. It is also well known that obesity can lead to or worsen SDB in both pediatric and adult populations [73]. This might be secondary to fat deposition in the neck causing narrow airway and/or a decrease in lung functional capacity due to truncal obesity. Obese children have a higher incidence of SDB and excessive daytime sleepiness (EDS) symptoms. This might be explained by the microarousals occurring at nighttime and oxygen desaturation, thus leading to EDS. It has been hypothesized that children with EDS compensate for their sleepiness by increased motor activity, thus presenting with what appear to be ADHD symptoms [74].

Given that SDB co-occurs with children's ADHD symptoms, a thorough clinical sleep evaluation should occur among children presenting with ADHD symptoms. Currently, many of these children are likely being treated for their ADHD symptoms with medication. It remains controversial as to whether treating associated OSA would cure or improve ADHD. Adenotonsillectomy, a commonly performed surgery today utilized to treat OSA, is relatively simple and improves up to 85% of children with SDB. Also, while ADHD symptoms may remain for some children, it is possible that improvement in SDB may allow for the use of lower medication doses to treat such residual ADHD symptoms.

Several limitations to the current meta-analysis, which incorporated a relatively large number of studies to estimate the ES between SDB and ADHD symptoms, deserve mention. Inclusion and exclusion criteria varied among different studies; e.g., some studies excluded individuals with a history of SDB or ADHD symptoms while others included them. Similarly, some studies included children with primary snoring in their SDB group, while others did not. Such study differences may affect the ES between SDB and ADHD symptoms. Furthermore, most studies focused on the use of clinical rating scales rather than diagnostic criteria per se, so it is unknown whether results would differ if data was based on diagnostic interviews or clinician diagnoses. Articles utilizing neuropsychological measures of attention were excluded as they generally correlate poorly with ADHD [75]. There are 21 studies that were excluded for this reason, most of which did not find a significant relation between SDB and ADHD symptoms [76]. Inclusion of these studies in the meta-analysis would have lowered the ES. Finally, most of the studies excluded most of the major medical and neurological problems. However, most of the studies did not directly comment on the comorbid psychiatric problems. Because included studies did not report on the presence of restless leg syndrome (RLS), a disorder with higher prevalence in children with ADHD, it is possible that the relationship between SDB and ADHD symptoms might be related to RLS symptoms and should be examined in future research [77].

Clinically, the present findings highlight the potential importance of screening children with ADHD symptoms for possible SDB symptoms. Of note, SDB is usually secondary to enlargement of the adenoids and the tonsils in most pediatric cases. This enlargement usually occurs between 2 and 8 years of age, coinciding with the typical onset of ADHD [78]. Thus, children diagnosed with ADHD symptoms in this age range might benefit from OSA/SDB screening. Such screening usually includes inquiries about snoring, observed apneas and waking up gasping for air. Oral examination shows enlarged tonsils, although adenoidal enlargement requires X-ray to facilitate diagnosis. Referral to a sleep specialist and PSG is required in cases with positive screening. AT is the treatment of choice with reassessment of the need for ADHD medication post-surgery.

Future research is needed to evaluate the ADHD symptom – SDB relation more extensively. For example, a better understanding of whether SDB is more closely related to ADHD symptoms or to

symptoms of sluggish cognitive tempo, which is often closely related to ADHD [79] or other psychiatric problems such as depression [70], is needed. More generally, prospective studies are needed to identify children and adolescents who are being treated for ADHD, screen them for SDB, and then reevaluate them for improvement of ADHD symptoms and further medication adjustment. We would then be able to truly evaluate the impact of treating co-morbid SDB in patients with ADHD and develop further guidelines for diagnosing and managing children with ADHD and associated SDB.

Practice points

- 1) Children and adolescent patients suffering from SDB present with elevated rates of ADHD symptoms. A significant improvement in ADHD symptoms occurred in children and adolescents after adenotonsillectomy compared to pre-surgery.
- 2) Children and adolescents presenting with ADHD symptoms should be screened for SDB and, if present, treated for their SDB before to initiating psychotropic medication.
- 3) Treating SDB in children and adolescents with ADHD symptoms might eliminate or decrease the need for psychotropic medication use.

Research agenda

- 1) Further research is needed to evaluate the etiological reasons for the relationship between SDB and ADHD symptoms.
- 2) In this meta-analysis, ADHD scales were used to assess ADHD symptoms. ADHD scales, while correlated with ADHD diagnosis, do not necessarily indicate a diagnosis of ADHD. Thus, research is needed to examine whether the relation with SDB holds for children clinically diagnosed with ADHD.
- 3) Prospective studies are needed to identify children and adolescents who are being treated for ADHD, screen them for SDB, and then reevaluate them for improvement of ADHD symptoms and further medication adjustment.

Conflict of interest

None.

Acknowledgment

None.

References*

- [1] Polanczyk G, de Lima MS, Horta BL, Biederman J, Rohde LA. The worldwide prevalence of ADHD: a systematic review and meta-regression analysis. *Am J Psychiatry* 2007;164:942–8.
- [2] American Psychiatric Association. Diagnostic criteria from DSM-IV-TR. Washington, D.C.: American Psychiatric Association; 2000.

* The most important references are denoted by an asterisk.

- [3] Reitman D, Hummel R, Franz DZ, Gross AM. A review of methods and instruments for assessing externalizing disorders: theoretical and practical considerations in rendering a diagnosis. *Clin Psychol Rev* 1998;18:555–84.
- [4] Sadeh A, Pergamin L, Bar-Haim Y. Sleep in children with attention-deficit hyperactivity disorder: a meta-analysis of polysomnographic studies. *Sleep Med Rev* 2006;10:381–98.
- [5] Revenagh PC, Chmielewski LJ, Edwards T, Krishna J, Krakovitz P, Anne S. Utility of preoperative cardiac evaluation in pediatric patients undergoing surgery for obstructive sleep apnea. *Arch Otolaryngol Head Neck Surg* 2011;137:1269–75.
- [6] Iber C, Ancoli-Israel S, Chesson A, Quan S. The AASM manual for the scoring of sleep and associated events: rules terminology, and technical specifications. Westchester, IL: Am Acad Sleep Med; 2007.
- [7] Sedky K, Carvalho K, Lippmann S. Attention deficit hyperactivity disorder and sleep disordered breathing in children. *J Pediatr Biochem* 2013;3:61–7.
- [8] Wei JL, Mayo MS, Smith HJ, Reese M, Weatherly RA. Improved behavior and sleep after adenotonsillectomy in children with sleep-disordered breathing. *Arch Otolaryngol Head Neck Surg* 2007;133:974–9.
- [9] Wei JL, Bond J, Mayo MS, Smith HJ, Reese M, Weatherly RA. Improved behavior and sleep after adenotonsillectomy in children with sleep-disordered breathing: long-term follow-up. *Arch Otolaryngol Head Neck Surg* 2009;135:642–6.
- [10] Gruber R, Xi T, Frenette S, Robert M, Vannasinh P, Carrier J. Sleep disturbances in prepubertal children with attention deficit hyperactivity disorder: a home polysomnography study. *Sleep* 2009;32:343–50.
- [11] Dillon JE, Blunden S, Ruzicka DL, Guire KE, Champagne D, Weatherly RA, et al. DSM-IV diagnoses and obstructive sleep apnea in children before and 1 year after adenotonsillectomy. *J Am Acad Child Adolesc Psychiatry* 2007;46:1425–36.
- *[12] Huang YS, Chen NH, Li HY, Wu YY, Chao CC, Guilleminault C. Sleep disorders in Taiwanese children with attention deficit/hyperactivity disorder. *J Sleep Res* 2004;13:269–77.
- *[13] O'Brien LM, Holbrook CR, Mervis CB, Klaus CJ, Bruner JL, Raffield TJ, et al. Sleep and neurobehavioral characteristics of 5- to 7-year-old children with parentally reported symptoms of attention-deficit/hyperactivity disorder. *Pediatrics* 2003;111:554–63.
- [14] Sangal RB, Owens JA, Sangal J. Patients with attention-deficit/hyperactivity disorder without observed apneic episodes in sleep or daytime sleepiness have normal sleep on polysomnography. *Sleep* 2005;28:1143–8.
- [15] Chervin RD, Archbold KH. Hyperactivity and polysomnographic findings in children evaluated for sleep-disordered breathing. *Sleep* 2001;24:313–20.
- *[16] Galland BC, Tripp EG, Gray A, Taylor BJ. Apnea-hypopnea indices and snoring in children diagnosed with ADHD: a matched case-control study. *Sleep Breath* 2011;15:455–62.
- [17] Cooper J, Tyler L, Wallace I, Burgess KR. No evidence of sleep apnea in children with attention deficit hyperactivity disorder. *Clin Pediatr* 2004;43:609–14.
- [18] Bass JL, Corwin M, Gozal D, Moore C, Nishida H, Parker S, et al. The effect of chronic or intermittent hypoxia on cognition in childhood: a review of the evidence. *Pediatrics* 2004;114:805–16.
- [19] Paavonen EJ, Porkka-Heiskanen T, Lähikainen AR. Sleep quality, duration and behavioral symptoms among 5–6-year-old children. *Eur Child Adolesc Psychiatry* 2009;18:747–54.
- [20] Li AM, Chan MH, Yin J, So HK, Cjam IH, Lam CW, et al. C-reactive protein in children with obstructive sleep apnea and the effects of treatment. *Pediatr Pulmonol* 2008;43:34–40.
- [21] Owens JA. Neurocognitive and behavioral impact of sleep disordered breathing in children. *Pediatr Pulmonol* 2009;44:417–22.
- [22] Huang YS, Guilleminault C, Li HY, Yang CM, Wu YY, Chen NH. Attention-deficit/hyperactivity disorder with obstructive sleep apnea: a treatment outcome study. *Sleep Med* 2007;8:18–30.
- [23] Chervin RD, Hedger K, Dillon JE, Pituch KJ. Pediatric sleep questionnaire (PSQ): validity and reliability of scales for sleep-disordered breathing, snoring, sleepiness, and behavioral problems. *Sleep Med* 2000;1:21–32.
- [24] Weber SA, Lima Neto AC, Ternes FJ, Montovani JC. Hyperactivity and attention deficit syndrome in obstructive sleep apnea syndrome: is there improvement with surgical management? *Braz J Otorhinolaryngol* 2006;72:124–9.
- [25] Chervin RD, Dillon JE, Bassetti C, Ganoczy DA, Pituch KJ. Symptoms of sleep disorders, inattention, and hyperactivity in children. *Sleep* 1997;20:1185–92.
- [26] Ali NJ, Pitson DJ, Stradling JR. Snoring, sleep disturbance, and behaviour in 4–5 year olds. *Arch Dis Child* 1993;68:360–6.
- [27] Ali NJ, Pitson D, Stradling JR. Natural history of snoring and related behaviour problems between the ages of 4 and 7 years. *Arch Dis Child* 1994;71:74–6.
- [28] Avior G, Fishman G, Leor A, Sivan Y, Kaysar N, Derowe A. The effect of tonsillectomy and adenoidectomy on inattention and impulsivity as measured by the Test of Variables of Attention (TOVA) in children with obstructive sleep apnea syndrome. *Otolaryngol Head Neck Surg* 2004;131:367–71.
- [29] Fidan TFV. The impact of adenotonsillectomy on attention-deficit hyperactivity and disruptive behavioral symptoms. *EAJM* 2008;40:14–7.
- *[30] Galland BC, Dawes PJ, Tripp EG, Taylor BJ. Changes in behavior and attentional capacity after adenotonsillectomy. *Pediatr Res* 2006;59:711–6.
- [31] Esteller More E, Barcelo Mongil M, Segarra Isern F, Pineiro Aguin Z, Pujol Olmo A, Soler EM, et al. Neurocognitive and behavioural abnormalities in paediatric sleep-related breathing disorders. *Acta Otorrinolaringol Esp* 2009;60:325–31.
- [32] Owens J, Spirito A, Marcotte A, McGuinn M, Berkelhammer L. Neuropsychological and behavioral correlates of obstructive sleep apnea syndrome in children: a preliminary study. *Sleep Breath* 2000;4:67–78.
- [33] Li HY, Huang YS, Chen NH, Fang TJ, Lee LA. Impact of adenotonsillectomy on behavior in children with sleep-disordered breathing. *Laryngoscope* 2006;116:1142–7.
- [34] Pagel JF, Snyder S, Dawson D. Obstructive sleep apnea in sleepy pediatric psychiatry clinic patients: polysomnographic and clinical correlates. *Sleep Breath* 2004;8:125–31.
- [35] Goraya JS, Cruz M, Valencia I, Kaleyias J, Khurana DS, Hardison HH, et al. Sleep study abnormalities in children with attention deficit hyperactivity disorder. *Pediatr Neurol* 2009;40:42–6.
- [36] Owens JA, Mehlenbeck R, Lee J, King MM. Effect of weight, sleep duration, and comorbid sleep disorders on behavioral outcomes in children with sleep-disordered breathing. *Arch Pediatr Adolesc Med* 2008;162:313–21.
- [37] Silvestri R, Gagliano A, Arico I, Calarese T, Cedro C, Bruni O, et al. Sleep disorders in children with Attention-Deficit/Hyperactivity Disorder (ADHD) recorded overnight by video-polysomnography. *Sleep Med* 2009;10:1132–8.
- [38] Melendres MC, Lutz JM, Rubin ED, Marcus CL. Daytime sleepiness and hyperactivity in children with suspected sleep-disordered breathing. *Pediatrics* 2004;114:768–75.
- [39] Brockmann PE, Urschitz MS, Schlaud M, Poets CF. Primary snoring in school children: prevalence and neurocognitive impairments. *Sleep Breath* 2012;16:23–9.
- *[40] Chervin RD, Ruzicka DL, Giordani BJ, et al. Sleep-disordered breathing, behavior, and cognition in children before and after adenotonsillectomy. *Pediatrics* 2006;117:e769–78.
- [41] Giordani B, Hodges EK, Guire KE, Ruzicka DL, Dillon JE, Weatherly RA, et al. Changes in neuropsychological and behavioral functioning in children with and without obstructive sleep apnea following tonsillectomy. *J Int Neuropsychol Soc* 2012;18:212–22.
- [42] Giordani B, Hodges EK, Guire KE, Ruzicka DL, Dillon JE, Weatherly RA, et al. Neuropsychological and behavioral functioning in children with and without obstructive sleep apnea referred for tonsillectomy. *J Int Neuropsychol Soc* 2008;14:571–81.
- [43] O'Brien LM, Ivanenko A, Crabtree VM, Holbrook CR, Bruner JL, Klaus CJ, et al. The effect of stimulants on sleep characteristics in children with attention deficit/hyperactivity disorder. *Sleep Med* 2003;4:309–16.
- [44] O'Brien LM, Ivanenko A, Crabtree VM, Holbrook CR, Bruner JL, Klaus CJ, et al. Sleep disturbances in children with attention deficit hyperactivity disorder. *Pediatr Res* 2003;54:237–43.
- [45] O'Brien LM, Mervis CB, Holbrook CR, Bruner JL, Smith NH, McNally N, et al. Neurobehavioral correlates of sleep-disordered breathing in children. *J Sleep Res* 2004;13:165–72.
- [46] Janusz JA. An examination of neuropsychological and behavioral functioning of children who snore. *Sci Eng* 1999.
- [47] Mulvaney S. Behavioral and cognitive correlates of sleep-disordered breathing in a community sample of school children; 2003.
- *[48] Kaemingk KL, Pasvogel AE, Goodwin JL, Mulvaney SA, Martinez F, Enright PL, et al. Learning in children and sleep disordered breathing: findings of the Tucson Children's Assessment of Sleep Apnea (tuCASA) prospective cohort study. *J Int Neuropsychol Soc* 2003;9:1016–26.
- *[49] Lewin DS, Rosen RC, England SJ, Dahl RE. Preliminary evidence of behavioral and cognitive sequelae of obstructive sleep apnea in children. *Sleep Med* 2002;3:5–13.
- *[50] Mitchell RB, Kelly J. Child behavior after adenotonsillectomy for obstructive sleep apnea syndrome. *Laryngoscope* 2005;115:2051–5.
- *[51] Mitchell RB, Kelly J. Long-term changes in behavior after adenotonsillectomy for obstructive sleep apnea syndrome in children. *Otolaryngol Head Neck Surg* 2006;134:374–8.
- *[52] Mitchell RB, Kelly J. Behavioral changes in children with mild sleep-disordered breathing or obstructive sleep apnea after adenotonsillectomy. *Laryngoscope* 2007;117:1685–8.
- [53] Mitchell RB, Boss EF. Pediatric obstructive sleep apnea in obese and normal-weight children: impact of adenotonsillectomy on quality-of-life and behavior. *Dev Neuropsychol* 2009;34:650–61.
- [54] Tran KD, Nguyen CD, Weedon J, Goldstein NA. Child behavior and quality of life in pediatric obstructive sleep apnea. *Arch Otolaryngol Head Neck Surg* 2005;131:52–7.
- [55] Chervin RD, Ruzicka DL, Hoban TF, Fetterolf JL, Garetz SL, Guire KE, et al. Esophageal pressures, polysomnography, and neurobehavioral outcomes of adenotonsillectomy in children. *Chest* 2012;142:101–10.
- [56] Blunden S, Lushington K, Kennedy D, Martin J, Dawson D. Behavior and neurocognitive performance in children aged 5–10 years who snore compared to controls. *J Clin Exp Neuropsychol* 2000;22:554–68.
- [57] Bourke RS, Anderson V, Yang JS, Jackman AR, Killedar A, Nixon GM, et al. Neurobehavioral function is impaired in children with all severities of sleep disordered breathing. *Sleep Med* 2011;12:222–9.
- [58] Gottlieb DJ, Chase C, Vezina RM, Heeren TC, Corwin MJ, Auerbach SH, et al. Sleep-disordered breathing symptoms are associated with poorer cognitive function in 5-year-old children. *J Pediatr* 2004;145:458–64.
- [59] Golan N, Shahar E, Ravid S, Pillar G. Sleep disorders and daytime sleepiness in children with attention-deficit/hyperactive disorder. *Sleep* 2004;27:261–6.
- [60] Landau YE, Bar-Yishay O, Greenberg-Dotan S, Goldbart AD, Tarasiuk A, Tal A. Impaired behavioral and neurocognitive function in preschool children with obstructive sleep apnea. *Pediatr Pulmonol* 2012;47:180–8.
- [61] Miano S, Paolino MC, Urbano A, Parisi P, Massolo AC, Castaldo R, et al. Neurocognitive assessment and sleep analysis in children with sleep-disordered breathing. *Clin Neurophysiol* 2011;122:311–9.

- [62] Rosen CL, Storfer-Isser A, Taylor HG, Kirchner HL, Emancipator JL, Redline S. Increased behavioral morbidity in school-aged children with sleep-disordered breathing. *Pediatrics* 2004;114:1640–8.
- [63] Ting H, Wong RH, Yang HJ, Lee SP, Lee SD, Wang L. Sleep-disordered breathing, behavior, and academic performance in Taiwan schoolchildren. *Sleep Breath* 2011;15:91–8.
- [64] Barnes ME, Gozal D, Molfese DL. Attention in children with obstructive sleep apnoea: an event-related potentials study. *Sleep Med* 2012;13:368–77.
- [65] Beebe DW, Ris MD, Kramer ME, Long E, Amin R. The association between sleep disordered breathing, academic grades, and cognitive and behavioral functioning among overweight subjects during middle to late childhood. *Sleep* 2010;33:1447–56.
- [66] Wells GASB, O'Connell D, Peterson J, Welch V, Losos M, Tugwell P. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analysis.
- [67] Cortese S, Faraone SV, Konofal E, Lecendreux M. Sleep in children with attention-deficit/hyperactivity disorder: meta-analysis of subjective and objective studies. *J Am Acad Child Adolesc Psychiatry* 2009;48:894–908.
- [68] Mulvaney SA, Goodwin JL, Morgan WJ, Rosen GR, Quan SF, Kaemingk KL. Behavior problems associated with sleep disordered breathing in school-aged children—the Tucson children's assessment of sleep apnea study. *J Pediatr Psychol* 2006;31:322–30.
- [69] Marcus CL, Moore RH, Rosen CL, Giordani B, Garetz CL, Taylor HG, et al. A randomized trial of adenotonsillectomy for childhood sleep apnea. *N Engl J Med* 2013;368:2366–76.
- [70] Yilmaz E, Sedky K, Bennett DS. The relationship between depressive symptoms and obstructive sleep apnea in pediatric populations: a meta-analysis. *J Clin Sleep Med* 2013;9:1213–20.
- [71] Feng J, Wu Q, Zhang D, Chen BY. Hippocampal impairments are associated with intermittent hypoxia of obstructive sleep apnea. *Chin Med J* 2012;125:696–701.
- [72] Mostofsky SH, Cooper KL, Kates WR, Denckla MB, Kaufmann WE. Smaller prefrontal and premotor volumes in boys with attention-deficit/hyperactivity disorder. *Biol Psychiatry* 2002;52:785–94.
- [73] Sedky K, Akhtar U, Oluwabusi O. The ABCDEs of obstructive sleep apnea. *Curr Psychiatry* 2013;12:2.
- [74] Cortese S, Konofal E, Dalla Bernardina B, Mouren MC, Lecendreux M. Does excessive daytime sleepiness contribute to explaining the association between obesity and ADHD symptoms? *Med Hypotheses* 2008;70:12–6.
- [75] Koziol LF, Stevens MC. Neuropsychological assessment and the paradox of ADHD. *Appl Neuropsychol Child* 2012;1:79–89.
- [76] Calhoun SL, Mayes SD, Vgontzas AN, Tsaousoglou M, Shifflett LJ, Bixler EO. No relationship between neurocognitive functioning and mild sleep disordered breathing in a community sample of children. *J Clin Sleep Med* 2009;5:228–34.
- [77] Miano S, Parisi P, Villa MP. The sleep phenotypes of attention deficit hyperactivity disorder: the role of arousal during sleep and implications for treatment. *Med Hypotheses* 2012;79:147–53.
- [78] Nanaware SK, Gothi D, Joshi JM. Sleep apnea. *Indian J Pediatr* 2006;73:597–601.
- [79] Barkley RA. Distinguishing sluggish cognitive tempo from ADHD in children and adolescents: executive functioning, impairment, and comorbidity. *J Clin Child Adolesc Psychol* 2013;42:161–73.